

Diagnostic Imaging Pathways - Seizure (Investigation)

Population Covered By The Guidance

This pathway provides guidance on the imaging of patients presenting with their first episode of seizure. A protocol for imaging patients with recalcitrant seizures is also included.

Date reviewed: August 2014

Date of next review: 2017/2018

Published: December 2014

Quick User Guide

Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.

Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

Pathway Diagram






Image Gallery

Note: These images open in a new page


1a Arterio-Venous Malformation

Image 1a and 1b (Magnetic Resonance Imaging): Axial and coronal images



demonstrating a large left parieto-occipital arterio-venous malformation (arrows).

1b



Teaching Points

- Patients without a provoked cause of first seizure require further evaluation through imaging
- In the emergency setting, contrast enhanced CT of the brain is a useful initial investigation
- MRI can depict subtle abnormalities that may go unrecognised on CT. Specific seizure protocols may be utilised in order to identify the epileptogenic focus
- Refractory epilepsy requires specialist referral. Video EEG, SPECT and PET have a role in this sub-group of patients

Computed Tomography (CT)

- When MRI is unavailable, CT is the next best imaging modality for establishing the cause of seizures and is able to detect early surgical lesions in the acute setting [11,12](#)
- Can identify large structural abnormalities and remains adequate in the emergency or perioperative setting [1,13,14](#)
- The role of contrast administration for both CT as well as MRI needs to be assessed [19](#)
- CT head with contrast enhancement may be useful in the setting of a focal seizure, neurological deficit, possible trauma, or absence of history of alcohol misuse
- Limitations: less sensitive than MRI at detecting the nature of the abnormality and demonstrating more subtle lesions such as encephalitis, hyperacute infarction, small mass lesions including tumours and vascular malformations, hippocampal sclerosis and developmental cortical malformations [7-10](#)

Magnetic Resonance Imaging (MRI)

- Imaging modality of choice for evaluation of epilepsy [1,2](#)
- Usually recommended in all patients with first seizure [3,4](#)
- Imaging technique used depends on the specific type of seizures suspected on clinical assessment and EEG eg a dedicated temporal lobe protocol for clinically suspected temporal lobe epilepsy or high resolution imaging through a possible epileptogenic region in those with clinically suspected extratemporal epilepsy [2](#)
- The identification of a lesion in extratemporal areas or of atrophy/increase signal in temporal lobe by qualitative or quantitative MRI, has a high correlation with the site of epileptogenesis [5](#)
- Allows diagnosis and provides prognostic information which can alter management for the

- individual patient with newly diagnosed partial seizures [6](#)
- Superior to CT in detection of cerebral lesions related to epilepsy, especially gliomas and cavernous malformations. Therefore, MRI may be needed to image patients with normal or inconclusive CT [3,7-10](#)
- MRI will be needed to detect and assess epileptogenic lesions unless there is an established provocative event
- Limitations
 - Limited availability and expense
 - Patients with contraindications to MRI

Magnetic Resonance Imaging (MRI) in Refractory Epilepsy

- Essential for pre-surgical evaluation of patients with uncontrolled epilepsy considered for surgery [15,16](#)
- MRI can provide quantitative and qualitative assessment and information and should be targeted to the area of clinical and EEG abnormalities (e.g. temporal or frontal lobe epilepsy) [1,5](#)
- Provides the anatomical resolution necessary for more accurate interpretation of functional imaging studies such as, SPECT and FDG-PET

Positron Emission Tomography (PET)

- Functional imaging technique allows presurgical localisation of seizure focus in patients with medically refractory partial seizures [5,18](#)
- During focal seizures, cerebral metabolic activity increases in the epileptogenic area, and PET reveals localised hypermetabolism. Interictal FDG-PET is useful for localisation of the epileptogenic region in patients with clinical syndrome of refractory mesial temporal lobe epilepsy or with suspected neocortical temporal lobe epilepsy [18](#)
- Interictal FDG-PET has sensitivity of 84% and specificity of 86% for temporal lobe epilepsy and 33% sensitivity and 95% specificity for extratemporal epilepsy [5](#)
- Useful in cases where there is discordant information eg EEG shows right temporal focus and MRI shows left mesial temporal sclerosis, or if MRI is normal in a patient with clinical/electrographic evidence of temporal epilepsy or if there is bilateral mesial temporal sclerosis. Also may be of value where there is "dual pathology" eg temporal sclerosis and a structural abnormality of uncertain significance elsewhere
- Limitations: expensive and limited availability

Single Photon Emission Computed Tomography (SPECT)

- Functional imaging technique useful in localisation of the seizure focus in the presurgical evaluation of patients with medically refractory epilepsy [5,17](#)
- Useful in cases where there is discordant data in temporal lobe epilepsy, non-lesional temporal or extratemporal epilepsy
- A seizure focus typically manifests as a focus of hypoperfusion on interictal examinations and as a focus of increased perfusion on ictal examinations [17](#)
- Ictal scans are compared with interictal baseline examination to detect subtle changes [17](#)
- Usually concurrent video EEG is required to allow correlation of the blood flow changes with clinical and electrophysiological changes

- Limitations: difficult interpretation requiring knowledge of seizure type, clinical activity, time of ictal injection in relation to seizure onset, and MRI findings [5](#)

References

References are graded from Level I to V according to the Oxford Centre for Evidence-Based Medicine, Levels of Evidence. [Download the document](#)

1. Connor SEJ, Jarosz JM. **Magnetic resonance imaging of patients with epilepsy.** Clin Radiol. 2001;56:787-801. (Review article)
2. Bradley WG, Shey RB. **MR imaging evaluation of seizures.** Radiology. 2000;214:651-6. (Review article)
3. King MA, Newton MR, Jackson GD, et al. **Epileptology of the first-seizure presentation: a clinical, electroencephalographic, and magnetic resonance imaging study of 300 consecutive patients.** Lancet. 1998;352:1007-11. (Level II evidence). [View the reference](#)
4. Scheuer ML, Pedley TA. **The evaluation and treatment of seizures.** N Engl J Med. 1990;323(21):1468-74. (Review article)
5. Spencer SS. **The relative contributions of MRI, SPECT, and PET imaging in epilepsy.** Epilepsia. 1994;35(S6):S72-S89. (Level III evidence)
6. Van Paesschen W, Duncan JS, Stevens JM, et al. **Etiology and early prognosis of newly diagnosed partial seizures in adults: a quantitative hippocampal MRI study.** Neurology. 1997;49:753-7. (Level III evidence)
7. Latack JT, Abou-Khalil BW, Siegel GJ, et al. **Patients with partial seizures: evaluation by MR, CT and PET imaging.** Radiology. 1986;159:159-63. (Level III evidence)
8. Laster DW, Penry JK, Moody DM, et al. **Chronic seizure disorders: contribution of MR imaging when CT is normal.** Am J Neuroradiol. 1985;6:177-80. (Level II/III evidence)
9. Bergen D, Bleck T, Ramsey R, et al. **Magnetic resonance imaging as a sensitive and specific predictor of neoplasms removed for intractable epilepsy.** Epilepsia. 1989;30:318-21. (Level III evidence)
10. Rigamonti D, Hadley MN, Drayer BP, et al. **Cerebral cavernous malformations: incidence and familial occurrence.** N Engl J Med. 1988;319:343-7. (Level III evidence)
11. Ramirez-Lassepas M, Cipolle RJ, Morillo LR, et al. **Value of computed tomographic scan in the evaluation of adult patients after their first seizure.** Ann Neurol. 1984;15:536-43. (Level III evidence)
12. Theodore WH, Dorwart R, Holmes M, et al. **Neuroimaging in refractory partial seizures: comparison of PET, CT and MRI.** Neurology. 1986;36:750-9. (Level III evidence)
13. Sempere AP, Villaverde FJ, Martinez-Menendez B, et al. **First seizure in adults: a prospective study from the emergency department.** Acta Neurol Scand. 1992;86:134-8. (Level III evidence)
14. Schoenenberger RA, Heim SM. **Indication for computed tomography of the brain in patients with first uncomplicated generalized seizure.** BMJ. 1994;309:986-9. (Level III evidence)
15. Commission on Neuroimaging of the International League Against Epilepsy. **Guidelines for neuroimaging evaluation of patients with uncontrolled epilepsy considered for surgery.** Epilepsia. 1998;39(2):1375-6. (Practice guideline)
16. Sperling MR, Wilson G, Engel J Jr, et al. **Magnetic resonance imaging in intractable partial epilepsy: correlative studies.** Ann Neurol. 1986;20:57-62. (Level II/III evidence) (Guideline document)
17. Mullan BP, O'Connor MK, Hung JC. **Single photon emission computed tomography.** Neuroimaging Clin N Am. 1995;5(4):647-73. (Review article)
18. DeCarli C, McIntosh AR, Blaxton TA. **Use of positron emission tomography for the evaluation of epilepsy.** Neuroimaging Clin N America. 1995;5(4):623-45. (Review article)



19. Harden CL, Huff JS, Schwartz TH, Dubinsky RM, Zimmerman RD, Weinstein S, Foltin JC, Theodore WH. **Reassessment: Neuroimaging in the emergency patient presenting with seizure (an evidence-based review): Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology.** *Neurology*. 2007;69:1772-80. (Level I evidence)

Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<p>Consent to Procedure or Treatment</p> <p>Radiation Risks of X-rays and Scans</p> <p>Computed Tomography (CT)</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Positron Emission Tomography (PET)</p>	<p>Computed Tomography (CT)</p> <p>Contrast Medium (Gadolinium versus Iodine)</p> <p>Gadolinium Contrast Medium</p> <p>Iodine-Containing Contrast Medium</p> <p>Magnetic Resonance Imaging (MRI)</p> <p>Plain Radiography/X-rays</p> <p>Radiation Risk of Medical Imaging During Pregnancy</p> <p>Radiation Risk of Medical Imaging for Adults and Children</p> <p>Nuclear Medicine</p> <p>PET Scan</p> <p>SPECT-CT Scan</p>

Copyright

© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change

File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.



[Legal Matters](#)