

# Diagnostic Imaging Pathways - Headache (Thunderclap)

## Population Covered By The Guidance

This pathway provides guidance on the imaging of adult patients with acute severe ('thunderclap') headache.

**Date reviewed: June 2014**

**Date of next review: 2017/2018**






**Published: October 2014**

## Quick User Guide

Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.


Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

## Pathway Diagram

Date reviewed: June 2014  
 Please note that this pathway is subject to review and revision




## Image Gallery

*Note: These images open in a new page*

1a


### Subarachnoid Haemorrhage

Image 1a and 1b (Computed Tomography): Pre and post-contrast images of a large left sided subarachnoid haemorrhage with blood surrounding the




Circle of Willis (Image 1a arrow) and extending into the ventricles (Image 1b arrow).

1b



1c



## Perimesencephalic Haemorrhage

- Perimesencephalic haemorrhage was first described in 1985 and describes a form of subarachnoid haemorrhage more benign in nature when compared to aneurysmal SAH [36](#)
- Approximately ten percent of spontaneous SAH are due to non aneurysmal perimesencephalic haemorrhage [37](#)
- In 95 percent of patients with a perimesencephalic pattern of haemorrhage in SAH the cause will be non-aneurysmal and this is an important clinical distinction as the prognosis for these patients is generally favourable. [38](#) The other 5 percent are typically due to vertebrobasilar aneurysm [39](#) with rarer causes including vascular tumours, arteriovenous malformations, dural arteriovenous fistula and trauma [37](#)
- Perimesencephalic haemorrhage is defined as the following [37](#)
  - Blood centered immediately anterior to the mid-brain or pons and may involve to differing degrees the interpeduncular, crural, ambient, quadrigeminal, prepontine or carotid cisterns
  - In addition to this, blood may thinly extend into the suprasellar cistern and the basal portions of the sylvian and interhemispheric fissures but not into the distal portions of the sylvian or interhemispheric fissures
  - There may be small amounts of blood that may sediment in the occipital horns of the lateral ventricles however there is no frank intraventricular haemorrhage

## Computed Tomography (CT) of the Head

Imaging modality of choice for investigation of thunderclap headache [1-3](#)

- High (greater than 90%) sensitivity for subarachnoid haemorrhage (SAH) if performed within 24 hours of haemorrhage [4-6](#)
- In a large multi center prospective cohort study, 953 patients who were neurologically intact and imaged within 6 hours of headache onset CT had a sensitivity of 100 percent and specificity of 100 percent. [7](#) Overall for 3132 patients enrolled the sensitivity of CT for SAH was 92.9 percent and specificity 100 percent. This has led further discussion as to whether CT can be considered as a 'rule out' for SAH. At this stage there remains conjecture over this proposal. The results acquired



- in this trial relied upon the availability of third generation CT scanners and dedicated neuroradiology expertise, logistical considerations that are not currently widely available
- Provides an estimate of the extent of the haemorrhage, and allows recognition of associated intracerebral haemorrhage, intraventricular haemorrhage, hydrocephalus, and risk of subsequent vasospasm [4](#)
  - A normal CT does not exclude subarachnoid haemorrhage
  - Limitations
    - Loss of sensitivity with increasing time between onset of headache and neuroimaging [4](#)
    - False negative results in cases of small-volume bleeding [4](#)
    - Possible false negative results for blood with a haematocrit of less than 30%
  - Cerebral venous thrombosis, another cause of thunderclap headache has the following CT features [8](#)
    - On pre-contrast CT the acute thrombus may be visible as an elongated high attenuation lesion within the dural sinus or cortical vein - '*the cord sign or dense triangle sign*'
    - On post-contrast images a filling defect may be seen as the dura enhances but the thrombus does not - '*the empty delta sign*'
    - CVT is found primarily in children and young adult and accounts for 1-2 percent of strokes [9](#)
  - Information on CT for consumers [InsideRadiology](#)

## Computed Tomography Angiography (CTA)

- Non-invasive imaging modality for demonstrating vascular anatomy, with greater than 90% sensitivity for depiction of cerebral aneurysms greater than 3mm in size [10](#)
- There are reports in the existing literature of CTA being able to detect active bleeding from aneurysms [11,12](#)
- Experience with CTA is still in its early stages and most patients will also have digital subtraction angiography (DSA) for definite diagnosis and management decisions. However, some neurosurgeons are using 3D-CT angiography in place of DSA in the diagnosis of ruptured aneurysms and for defining vascular anatomy before surgery [10,13-15](#)
- Advances in imaging technology have led to the improved diagnostic accuracy of CTA. In a 2012 study [16](#) CTA had a sensitivity of 97.8 percent and specificity of 88.7 percent for the detection of intracranial aneurysms when compared against 3D DSA. Importantly these figures included the detection of aneurysms less than 3mm, a previous limitation with earlier technology. The results of this trial were concordant with other trials investigating these parameters. [16-21](#) A 2012 systematic review and meta-analysis found the pooled sensitivity and specificity of CT Angiography for detected cerebral aneurysms in acute SAH as 98 percent and 100 percent respectively [22](#)
- Limitations
  - Limited evaluation of collateral pathways and cerebral haemodynamics
  - Inferior spatial resolution compared to DSA
  - Does not provide precise information about intracranial haemodynamics
  - Does not offer therapeutic opportunity
- Information on CT for consumers [InsideRadiology](#)

## Digital Subtraction Angiography (DSA)

- Digital subtraction angiography is a procedure where a control radiograph is taken prior to contrast. Contrast is administered into a blood vessel and a radiograph is again taken, before sophisticated computer software highlights the vascular structure with contrast and removes

- surrounding tissues
- Gold standard for the detection of ruptured intracranial aneurysms and depicting the cerebral vascular anatomy [3,23](#)
- Highly sensitive for detection of small aneurysms, small arteriovenous malformations and dural vascular malformations [23](#)
- SAH is found to have no vascular origin with initial catheter angiography in approximately 15 percent of cases [24](#)
- If the initial DSA is normal and the pattern of haemorrhage on original CT is perimesencephalic in distribution there is generally considered no need for a repeat angiogram [13,14](#)
- For patients with an initial negative DSA and where repeat DSA is conducted, a 2013 study found that in the subsequent instances a cause of SAH was found in 4.5 percent of cases reviewed. [27](#) A similar trial analysing the utility of repeat DSA in patients with perimesencephalic subarachnoid haemorrhage had similar results with 3.3 percent of cases demonstrating a causative vascular lesion [27](#) though in a larger study the diagnostic yield of repeat angiography in patients with perimesencephalic haemorrhage was 0 percent [24](#)
- Advantages
  - Offers therapeutic opportunity
  - Better outcome in terms of disability at 1 year with endovascular coiling compared to neurosurgical clipping [28](#)
- Disadvantages
  - Invasive procedure with potential complications

## Lumbar Puncture

- Performed in cases of suspected subarachnoid haemorrhage (SAH) with negative, equivocal or technically inadequate CT [5,6,29](#)
- If SAH is suspected, the lumbar puncture should be performed at least 6, and preferably 12 hours CT after the onset of suspected SAH [30](#)
- CSF should be examined for xanthochromia by means of spectrophotometry [30](#)
- If CT or lumbar puncture indicates the presence of SAH, digital subtraction angiography is indicated to identify the cause [1](#)
- If the CT and lumbar puncture are both normal subarachnoid haemorrhage has been effectively excluded [1,3,32](#)
- Limitations [7](#)
  - Invasive with potential exacerbation of symptoms
  - Possible misinterpretation of 'traumatic tap' as SAH

## Magnetic Resonance Imaging (MRI) / MR Angiography (MRA) / MR Venography (MRV)

- The combination of MRI and magnetic resonance venography (MRV) is the imaging modality of choice for the investigation of suspected cerebral venous thrombosis (CVT) [33](#)
- Cerebral venous thrombosis has the following features [33](#)
  - Highly variable and non specific presentation from thunderclap headache to symptoms of raised cerebral venous pressure such as headache, vomiting and papilloedema
  - Imaging findings can be direct when the thrombus is visible within the cerebral venous system or indirect when there are ischaemic changes visible related to the venous outflow obstruction [34](#)



- On MRI acute thrombus is isointense to brain on T1-weighted images and hypointense on T2-weighted images. Between 3 and 7 days after thrombus formation the clot becomes hyperintense on T-1 weighted images [8](#)
- MRI is also sensitive to the parenchyma and haemorrhagic changes of venous infarction [35](#)
- Information on Magnetic Resonance Imaging for consumers [InsideRadiology](#)

## References

Date of literature search: June 2014

The search methodology is available on request. [Email](#)

References are graded from Level I to V according to the Oxford Centre for Evidence-Based Medicine, Levels of Evidence. [Download the document](#)

- 1.** Edlow JA, Panagos PD, Godwin SA, Thomas TL, Decker WW. **Clinical policy: critical issues in the evaluation and management of patients presenting to the emergency department with acute headache.** Ann Emerg Med. 2008;52(4):407-36. (Review article)
- 2.** Field AG, Wang E. **Evaluation of the nontraumatic headache: an evidence based approach.** Emerg Med Clin North Am. 1999;17(1):127-52. (Review article)
- 3.** Mayberg MR, Batjer HH, Dacey R, et al. **Guidelines for the management of aneurysmal subarachnoid haemorrhage: a statement for healthcare professionals from a special writing group of the stroke council, American Heart Association.** Stroke. 1994;25(11):2315-28. (Review article)
- 4.** Kassell NF, Torner JC, Haley EC Jr, et al. **The international cooperative study on the timing of aneurysm surgery, part 1: overall management results.** J Neurosurg. 1990;73:18-36. (Level II evidence). [View the reference](#)
- 5.** Van der Wee N, Rinkel GJ, Hasan D, et al. **Detection of subarachnoid haemorrhage on early CT: is lumbar puncture still needed after a negative scan?** J Neurol Neurosurg Psychiatry. 1995;58:357-9. (Level II evidence). [View the reference](#)
- 6.** Morgenstern LB, Luna-Gonzales H, Huber JC Jr, et al. **Worst headache and subarachnoid hemorrhage: Prospective, Modern computed tomography and spinal fluid analysis.** Ann Emerg Med. 1998;32:297-304. (Level II evidence). [View the reference](#)
- 7.** Perry JJ, Stiell IG, Sivillotti ML, Bullard MJ, Emond M, Symington C, et al. **Sensitivity of computed tomography performed within six hours of onset of headache for diagnosis of subarachnoid haemorrhage: prospective cohort study.** BMJ. 2011;343:d4277. (Level I evidence)
- 8.** Lee SK, terBrugge KG. **Cerebral venous thrombosis in adults: the role of imaging evaluation and management.** Neuroimaging Clin N Am. 2003;13(1):139-52. (Review article)
- 9.** Sahin N, Solak A, Genc B, Bilgic N. **Cerebral venous thrombosis as a rare cause of subarachnoid hemorrhage: case report and literature review.** Clin Imaging. 2014;38(4):373-9. (Review article)
- 10.** Anderson GB, Steinke DE, Petruk KC, Ashforth R, Findlay JM. **Computed tomographic angiography versus digital subtraction angiography for the diagnosis and early treatment of ruptured intracranial aneurysms.** Neurosurgery. 1999;45(6):1315-20; discussion 20-2. (Level II evidence)
- 11.** Kathuria S, Deveikis JP, Westesson PL, Gandhi D. **Improved diagnosis of actively bleeding aneurysm on CT angiography using delayed CT images.** Eur J Radiol. 2011;79(2):328-31. (Review article)
- 12.** Desai S, Friedman JA, Hlavin J, Kash F. **Actively bleeding intracranial aneurysm demonstrated by CT angiography.** Clin Neurol Neurosurg. 2009;111(1):94-6. (Review article)



13. Hashimoto H, Iida J, Hironaka Y, Okada M, Sakaki T. **Use of spiral computerized tomography angiography in patients with subarachnoid hemorrhage in whom subtraction angiography did not reveal cerebral aneurysms.** J Neurosurg. 2000;92(2):278-83. (Level II evidence)
14. Matsumoto M, Sato M, Nakano M, Endo Y, Watanabe Y, Sasaki T, et al. **Three-dimensional computerized tomography angiography-guided surgery of acutely ruptured cerebral aneurysms.** J Neurosurg. 2001;94(5):718-27. (Level II evidence)
15. Velthuis BK, van Leeuwen MS, Witkamp TD, Ramos LM, Berkelbach van der Sprenkel JW, Rinkel GJ. **Surgical anatomy of the cerebral arteries in patients with subarachnoid hemorrhage: comparison of computerized tomography angiography and digital subtraction angiography.** J Neurosurg. 2001;95(2):206-12. (Level II evidence)
16. Lu L, Zhang LJ, Poon CS, Wu SY, Zhou CS, Luo S, et al. **Digital subtraction CT angiography for detection of intracranial aneurysms: comparison with three-dimensional digital subtraction angiography.** Radiology. 2012;262(2):605-12. (Level II evidence)
17. Donmez H, Serifov E, Kahriman G, Mavili E, Durak AC, Menku A. **Comparison of 16-row multislice CT angiography with conventional angiography for detection and evaluation of intracranial aneurysms.** Eur J Radiol. 2011;80(2):455-61. (Level II evidence)
18. He W, Hauptman J, Pasupuleti L, Setton A, Farrow MG, Kasper L, et al. **True posterior communicating artery aneurysms: are they more prone to rupture? A biomorphometric analysis.** J Neurosurg. 2010;112(3):611-5. (Level II evidence)
19. Struffert T, Doelken M, Adamek E, Schwarz M, Engelhorn T, Kloska S, et al. **Flat-detector computed tomography with intravenous contrast material application in experimental aneurysms: comparison with multislice CT and conventional angiography.** Acta Radiol. 2010;51(4):431-7. (Level II evidence)
20. Tipper G, JM UK-I, Price SJ, Trivedi RA, Cross JJ, Higgins NJ, et al. **Detection and evaluation of intracranial aneurysms with 16-row multislice CT angiography.** Clin Radiol. 2005;60(5):565-72. (Level II evidence)
21. Yoon DY, Lim KJ, Choi CS, Cho BM, Oh SM, Chang SK. **Detection and characterization of intracranial aneurysms with 16-channel multidetector row CT angiography: a prospective comparison of volume-rendered images and digital subtraction angiography.** AJNR Am J Neuroradiol. 2007;28(1):60-7. (Level II evidence). [View the reference](#)
22. Westerlaan HE, van Dijk JM, Jansen-van der Weide MC, de Groot JC, Groen RJ, Mooij JJ, et al. **Intracranial aneurysms in patients with subarachnoid hemorrhage: CT angiography as a primary examination tool for diagnosis - systematic review and meta-analysis.** Radiology. 2011;258(1):134-45. (Level I evidence). [View the reference](#)
23. White PM, Wardlaw JM, Easton V. **Can noninvasive imaging accurately depict intracranial aneurysms? A systematic review.** Radiology. 2000;217(2):361-70. (Level II evidence). [View the reference](#)
24. Rinkel GJ, Wijndicks EF, Hasan D et al. **Outcome in patients with subarachnoid hemorrhage and negative angiography according to pattern of haemorrhage on computed tomography.** Lancet. 1991;338:964-8. (Level II evidence). [View the reference](#)
25. Rinkel GJ, Wijndicks EF, Vermeulen M et al. **The clinical course of perimesencephalic subarachnoid haemorrhage.** Ann Neurol. 1991;29:463-8. (Level II evidence). [View the reference](#)
26. Ringelstein A, Mueller O, Goericke SL, Moeninghoff C, Sure U, Wanke I, et al. **Benefit of Second Catheter Angiography in Patients with Nontraumatic Subarachnoidal Hemorrhage.** Clin Neuroradiol. 2013:(Ahead of print - Level III evidence)
27. Molyneux A, Kerr R, Stratton I, Sandercock P, Clarke M, Shrimpton J, et al. **International Subarachnoid Aneurysm Trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised trial.** Lancet. 2002;360(9342):1267-74. (Level I evidence)
28. Edlow JA, Wyer PC. **Evidence-based emergency medicine/clinical question. How good is a negative cranial computed tomographic scan result in excluding subarachnoid**



**hemorrhage?** Ann Emerg Med. 2000;36(5):507-16. (Review article)

29. Vermeulen M, van Gijn J. **The diagnosis of subarachnoid haemorrhage.** J Neurol Neurosurg Psychiatry. 1990;53:365-72. (Review article)

30. Wijdicks EFM, Kerkhoff H, van Gijn J. **Long-term follow-up of 71 patients with thunderclap headache mimicking subarachnoid haemorrhage.** Lancet. 1988;2:68-70. (Level III evidence)

31. Harling DW, Peatfield RC, Van Hille PT, Abbott RJ. **Thunderclap headache: is it migraine?** Cephalalgia. 1989;9:87-90. (Level III evidence)

32. Tsai FY, Wang AM, Matovich VB, et al. **MR staging of acute dural sinus thrombosis: correlation with venous pressure measurements and implications for treatment and prognosis.** AJNR Am J Neuroradiol. 1995;16:1021-9. (Level III evidence)

33. Lee SK, terBrugge KG. **Cerebral venous thrombosis in adults: the role of imaging evaluation and management.** Neuroimaging Clin N Am. 2003;13:139-52. (Review article)

34. Forbes KP, Pipe JG, Heiserman JE. **Evidence for cytotoxic edema in the pathogenesis of cerebral venous infarction.** AJNR Am J Neuroradiol. 2001;22:450-5. (Level III evidence)

35. van Gijn J, van Dongen KJ, Vermeulen M, Hijdra A. **Perimesencephalic hemorrhage: a nonaneurysmal and benign form of subarachnoid hemorrhage.** Neurology. 1985;35(4):493-7. (Level III evidence)

36. Marder CP, Narla V, Fink JR, Tozer Fink KR. **Subarachnoid hemorrhage: beyond aneurysms.** AJR Am J Roentgenol. 2014;202(1):25-37. (Review article). [View the reference](#)

37. Velthuis BK, Rinkel GJ, Ramos LM, Witkamp TD, van Leeuwen MS. **Perimesencephalic hemorrhage. Exclusion of vertebrobasilar aneurysms with CT angiography.** Stroke. 1999;30(5):1103-9. (Level II evidence). [View the reference](#)

38. van der Schaaf IC, Velthuis BK, Gouw A, Rinkel GJ. **Venous drainage in perimesencephalic hemorrhage.** Stroke. 2004;35(7):1614-8. (Level III evidence). [View the reference](#)

## Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<p><a href="#">Consent to Procedure or Treatment</a></p> <p><a href="#">Radiation Risks of X-rays and Scans</a></p> <p><a href="#">Angiography (Angiogram)</a></p> <p><a href="#">Computed Tomography (CT)</a></p> <p><a href="#">Computed Tomography (CT) Angiography</a></p> <p><a href="#">Magnetic Resonance Angiography (MRA)</a></p> <p><a href="#">Magnetic Resonance Imaging (MRI)</a></p>	<p><a href="#">Angiography</a></p> <p><a href="#">Computed Tomography (CT)</a></p> <p><a href="#">Contrast Medium (Gadolinium versus Iodine)</a></p> <p><a href="#">Gadolinium Contrast Medium</a></p> <p><a href="#">Iodine-Containing Contrast Medium</a></p> <p><a href="#">Magnetic Resonance Imaging (MRI)</a></p> <p><a href="#">Plain Radiography/X-rays</a></p> <p><a href="#">Radiation Risk of Medical Imaging During</a></p>





	<p><a href="#">Pregnancy</a></p> <p><a href="#">Radiation Risk of Medical Imaging for Adults and Children</a></p> <p><a href="#">Interventional Radiological Treatment of Intracranial Aneurysms</a></p> <p><a href="#">SAH Vasospasm Endovascular Treatment</a></p>
--	--

## Copyright


© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

## Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change

## File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.



## [Legal Matters](#)